

FETAL DEVELOPMENT OF THE LONG BONES OF LOWLAND PACA (*CUNICULUS PACA*) BY ULTRASONOGRAPHY

Gisela Bujaldón Carbó
Final Degree Project – June 2019

UAB
Universitat Autònoma
de Barcelona



Introduction

The paca (*Cuniculus paca*) is a frugivorous hystricomorphic rodent in the tropical forests of South and Central America. This species is appreciated for its meat, which makes it an important resource of subsistence for Amazon people, and has important ecological roles, as seed disperser and prey specie.

Material & Methods

- 101 paca embryos/fetuses collected from two locations in the Amazon forest (2002 - 2016).
- External biometrics measurement:** total dorsal length (TDL).
- Ultrasonography analysis:** measurement of each mineralized (hyperechoic) and non-mineralized (anechoic) structure in the thoracic limb (humerus and radius) and pelvic limb (femur and tibia).

Discussion

The onset of the diaphysis mineralization occurs earlier in hystricomorphs (34% of the gestational period) than in the lab rat (75% of the gestational period).

Compared with other altricial species (cat and dog), the newborn paca has all secondary ossification centers mineralized, showing earlier motor ability.




Figure 1. Ultrasound images showing the evolution of the femur in different fetal stages in lowland paca fetuses (*Cuniculus paca*). Longitudinal cut. **A.** Fetus with a TDL of 6,9 cm (bar: 5 mm). **B.** Fetus with a TDL of 10,4 cm (bar: 5 mm). **C.** Fetus with a TDL of 14 cm (bar: 9 mm). **D.** Fetus with a TDL of 16,5 cm (bar: 10 mm). **E.** Fetus with a TDL of 20,4 cm (bar: 10 mm). **F.** Fetus with a TDL of 22,2 cm (bar: 10 mm). **G.** Fetus with a TDL of 27,1 cm (bar: 10 mm). **H.** Fetus with a TDL of 35,9 cm (bar: 13 mm). Mineralized diaphysis (solid line); epiphysis (*); Distal epiphyseal nucleus (+).

Intrauterine bony development seems to be related to the habits this rodent will have during its postnatal life.

- Digging** → radius and distal humerus more robust.
- Swimming** → tibia larger and more robust than femur.




Figure 2. Relationship between total dorsal length (TDL) in embryos / fetuses of lowland paca (*Cuniculus paca*) and mineralized structures of the femur. **A.** Diaphysis length. **B.** Head nucleus diameter. **C.** Greater trochanter nucleus diameter. **D.** Distal nucleus diameter.




Figure 3. Probability curves for the occurrence of skeletal bone mineralization in 101 embryos/fetuses of lowland paca (*Cuniculus paca*) in relation to TDL.

Conclusions

Paca fetuses have a skeleton sufficiently developed to favor the newborn locomotion, and earlier compared to that in domestic carnivores.

References

- Biknevicius AR. 1993. Biomechanical Scaling of Limb Bones and Differential Limb Use in Caviomorph Rodents. *Journal of Mammalogy*. 74(1):95–107.
- El Bizri HR, Monteiro FOB, de Andrade R dos S, Valsecchi J, Guimarães DA de A, Mayor P. 2017. Embryonic and fetal morphology in the lowland paca (*Cuniculus paca*): A precocial hystricomorph rodent. *Theriogenology*. 104:7–17.
- Patton JT, Kaufman MH. 1995. The timing of ossification of the limb bones, and growth rates of various long bones of the fore and hind limbs of the prenatal and early postnatal laboratory mouse. *Journal of anatomy*. 186 (Pt 1):175–85.
- Samuels JX, Van Valkenburgh B. 2008. Skeletal indicators of locomotor adaptations in living and extinct rodents. *Journal of Morphology*. 269(11):1387–1411.